

Editorial

Viral regulation of cell death

Matthew Reeves

UCL Institute of Immunity & Transplantation, Royal Free Hospital, London, UK

Correspondence should be addressed to Matthew Reeves; Tel: +44 207 794 0500 ext 33109, Fax: +44 207 830 2854, E-mail: matthew.reeves@ucl.ac.uk

The cell autonomous immune response is multifaceted and forms an important first line of defence upon pathogen infection (Randow *et al.* 2013). A major element of this response is the induction of cell suicide pathways, which contribute to the limiting of infection through elimination of a replication niche. Consequently, pathogens have evolved a multitude of strategies to intercept this pathway to promulgate infection. Human Cytomegalovirus (HCMV), a human herpes virus, is no exception and expresses an armoury of gene functions that promote viability (McCormick 2008). However, a key biological characteristic of HCMV is the ability to establish lifelong latent infections which are hallmarked by limited viral gene expression and no viral replication (Dupont & Reeves 2016).

One of the first host cell 'trip wires' is the detection of the pathogen upon entry into the cell. Importantly, we and others have shown that the virus must ameliorate this response for the establishment of non-lytic but functional infections (Chan *et al.* 2010, Reeves *et al.* 2012). Crucially, the absence of many viral anti-apoptotic functions during this phase of infection has rendered the virus dependent on the activation of key cellular functions. The favoured model proposes that HCMV engages with the cell and triggers the activation of cell death pathways but, via concomitant activation of the survival pathways, the outcome is positive for the virus, resulting in long term infection of these cells.

For HCMV, and the establishment of latency in CD34+ cells, the activation of ERK-MAPK signalling appears to be a crucial event (Reeves *et al.* 2012). In the current issue, Kew *et al.* (2017) have investigated this in more detail and demonstrate that ERK activation underpins a multi-faceted survival response encompassing post translational down-regulation of pro-apoptotic Bim and the induction ELK-1 transcriptional activity. Importantly, the activation of ERK is required to prevent the consequence of a concomitant activation of the pro-apoptotic protein, Bak, by viral infection. The challenge now is to fully

understand how these multiple events tip the balance in favour of survival in the infected cells. For instance, deletion of ELK-1 was not overly deleterious to CD34+ cells until the cells were subsequently stressed by viral infection, supporting a role in acute survival responses rather than long term homeostasis of viability.

The question remains as to why this is important. Clearly, the immediacy of understanding how pathogens promote cell survival has direct impact on our understanding of the complexity of the host-pathogen interaction. However, the wider implications are driven by a desire to decipher the biology of the cell. Dysregulated cell death underpins the pathogenesis of many infectious and non-infectious diseases and thus, understanding how these pathways are regulated can give new insight and, potentially, new hope for treatment of many complex diseases. For example, a proof of principle study showed that a CMV-encoded RNA that maintained energy production in virally infected cells (Reeves *et al.* 2007) could be used to protect rodents from the symptoms of Parkinson's disease (Kuan *et al.* 2012). Ultimately, viruses are assiduous molecular biologists - experts at hijacking, partitioning or re-directing cellular functions. It is hoped that, as we further unravel and define the complexity of the host-pathogen interaction, we will provide the foundation to interrogate host cell biology and reveal the many secrets still hidden by Mother Nature.

References

- Chan G, Nogalski MT, Bentz GL, Smith MS, Parmater A & Yurochko AD 2010 PI3K-dependent upregulation of Mcl-1 by human cytomegalovirus is mediated by epidermal growth factor receptor and inhibits apoptosis in short-lived monocytes. *J Immunol* **184** 3213-3222
- Dupont L & Reeves MB 2016 Cytomegalovirus latency and reactivation: recent insights into an age old problem. *Rev Med Virol* **26** 75-89
- Kew V, Wills M & Reeves M 2017 HCMV activation

of ERK-MAPK drives a multi-factorial response promoting the survival of infected myeloid progenitors. *J Mol Biochem* **6** 13-25

Kuan WL, Poole E, Fletcher M, Karniely S, Tyers P, Wills M, Barker RA & Sinclair JH 2012 A novel neuroprotective therapy for Parkinson's disease using a viral noncoding RNA that protects mitochondrial complex I activity. *J Exp Med* **209** 1-10

Randow F, MacMicking JD & James LC 2013 Cellular self-defense: how cell-autonomous immunity protects against pathogens. *Science* **340** 701-706

McCormick AL 2008 Control of apoptosis by human cytomegalovirus. *Curr Top Microbiol Immunol* **325** 281-295

Reeves MB, Breidenstein A & Compton T 2012 Human cytomegalovirus activation of ERK and myeloid cell leukemia-1 protein correlates with survival of latently infected cells. *Proc Natl Acad Sci U S A* **109** 588-593

Reeves MB, Davies AA, McSharry BP, Wilkinson GW & Sinclair JH 2007 Complex I binding by a virally encoded RNA regulates mitochondria-induced cell death. *Science* **316** 1345-1348